

The effect of saccadic adaptation on localization of visual targets

Holger Awater^{1,4}, David Burr^{2,3}, Markus Lappe^{1,4},

M. Concetta Morrone^{2,5} and Michael E. Goldberg^{6,7}

¹Lehrstuhl für Allgemeine Zoologie und Neurobiologie, Ruhr Universität, Bochum D-44780, Germany

²Istituto di Neuroscienze del CNR, Pisa 56100, Italy

³Dipartimento di Psicologia, Università di Firenze, Italy.

⁴Abteilung für Psychologie, Münster Universität, Germany.

⁵Facoltà di Psicologia, Università Vita-Salute San Raffaele, Milano 20132, Italy.

⁶Mahoney Center for Brain and Behavior, Center for Neurobiology and Behavior, and Departments of Neurology and Psychiatry, Columbia University

⁷New York State Psychiatric Institute.

Running title: *Saccadic Adaptation and Visual Perception*

Correspondence should be addressed to:

Michael E. Goldberg

1051 Riverside Drive, Unit 87, New York, NY 10027

meg2008@columbia.edu

Voice: 212-543-6931 Extension 101

Fax 212-543-5816

ABSTRACT

Objects flashed briefly around the time of a saccadic eye-movement are grossly mislocalized by human subjects, so they appear to be compressed towards the endpoint of the saccade. In this study we investigate spatial localization during saccadic adaptation, to examine whether the focus of compression tends towards the intended saccadic target, or at the endpoint of the actual (adapted) movement. We report two major results. Firstly, that peri-saccadic focus of the compression does not occur at the site of the initial saccadic target, but tended towards the actual landing site of the saccade. Secondly, and more surprisingly, we observed a large long-term perceptual distortion of space, lasting for hundreds of milliseconds. This distortion did not occur over the whole visual field but was limited to a local region of visual space around the saccade target, suggesting that saccadic adaptation induces a visuo-topic remapping of space. The results imply that the mechanisms controlling saccadic adaptation also affect perception of space and point to a strong perceptual plasticity coordinated with the well-documented plasticity of the motor system.

INTRODUCTION

Human spatial visual perception is ordinarily veridical and accurate. However, humans dramatically mislocalize objects flashed briefly just before or during saccades, the rapid eye movement made to shift the point of gaze from one location to another. Mislocalizations in the dark have been reported to occur in the direction of the saccade, and are roughly proportional to saccadic size (Cai et al. 1997; Dassonville et al. 1992, 1995; Honda 1989; Matin and Pearce 1965). However, during conditions of normal room lighting, briefly-flashed peri-saccadic targets are mislocalized towards the endpoint of the saccade: stimuli flashed either left of right of the saccadic target are seen at the site of the target, implying a transient compression of visual space for briefly presented stimuli (Lappe et al. 2000; Morrone et al. 1997; Ross et al. 1997; Ross et al. 2001). The compression follows a precise time course, maximal at saccadic onset, stretching about 50 ms before and after. Compression has also been reported in total darkness, but it is less obvious and is related to a mislocalization of both the saccade target and the flashed object (Awater et al. 2000). Experiments with simulated saccades, using a moving mirror to duplicate the retinal events of a saccade without requiring that the subject actually make a saccade, clearly suggest that the apparent displacement of absolute position is mediated at least partially by an extraretinal signal, or corollary discharge (Morrone et al. 1997; see also Ross et al. 2001).

The saccadic system is highly adaptive, and can rapidly adjust to changes in the oculomotor musculature and plant that can be evoked by growth, injury (Kommerell et al. 1976) and fatigue (Barash et al. 1999). Although the adaptation occurs in the motor rather than perceptual system, the signal driving this adaptive process is the visual saccadic error, the distance between the fovea and the target at the end of the saccade (Wallman and Fuchs 1998). In the laboratory this saccadic plasticity can be mimicked by a technique called short-term saccadic adaptation (McLaughlin 1967), that takes advantage of the inability of human subjects to detect small target displacements during saccades (Bridgeman et al. 1975). During adaptation the saccade target steps in a stereotypical way at the beginning of the eye movement, towards or away from the fixation point. Although this target step is imperceptible to the subject, the oculomotor system interprets the saccadic error as being caused by an inappropriate saccade, and adjusts the amplitude of the saccade appropriately. The adjustment is gradual, so that after a number of trials the eye ultimately lands on the final position of the target rather than its initial position.

Although saccadic adaptation is thought to affect primarily the motor system (Hopp and Fuchs 2002; Wallman and Fuchs 1998), two studies have shown that it also has perception consequences for targets presented well before saccadic onset. Moidell and Bedell (1988) reported small (about 0.5°) but consistent errors in perceptual localization during a fixation task after prolonged saccadic adaptation, suggesting a generalized distortion of visual space. More recently, Bahcall and Kowler (1999) measured the perceptual consequences of adaptation more directly, by asking subjects to compare the apparent positions of pre- and post-saccadic probes during saccadic adaptation. The results show that subjects perceived the post-saccadic probes at the position where the eyes actually moved, even though this was quite far from the target that elicited the saccade. Bahcall and Kowler interpreted these results to imply that the “efference copy” signal used to stabilize perception during saccades is based on the intended saccade rather than the actual eye-movement.

As mentioned previously, studies of peri-saccadic mislocalization show that the mislocalization is more complex than a simple shift, depending critically on retinal location: over a wide range of retinal positions, stimuli are localized towards a very strong attractor, the saccadic target. In this study we plan to take advantage of this phenomenon to understand better the nature of the efference copy signal during saccadic adaptation, where the intended saccadic target is different from the actual target. To this end we measured visual mislocalization of brief stimuli presented before, during and after saccades, during conditions of saccadic adaptation. In particular we were interested in whether the focus of peri-saccadic compression would be determined by an intention-to-move signal, or by the motor signal driving the actual saccade.

We report two major results. Firstly, that peri-saccadic focus of the compression (Ross et al. 1997) does not occur at the site of the initial saccadic target, but tended towards the actual landing site of the saccade. Secondly, and more surprisingly, we observed a large long-term perceptual distortion of space, lasting for hundreds of milliseconds. This distortion did not occur over the whole visual field but was limited to a local region of visual space around the saccade target, suggesting that saccadic adaptation induces a visuo-topic remapping of space. An abstract of these results has been presented elsewhere (Awater et al. 2001).

METHODS

Visual stimuli: Stimuli were displayed on a Barco color monitor with display area 36 x 25 cm, subtending 60 x 40° at the viewing distance of 37 cm. The display was surrounded by the dark brown frame of the monitor, lit to about 1 cd/m² by the dim background illumination. Stimuli were generated at 250 Hz by a visual stimulus generator (Cambridge Research Systems VSG2) housed in a personal computer (PC), programmed in C. All stimuli were presented on a red background (Commission Internationale de l'Eclairage (CIE) coordinates: $x=0.595$; $y=0.349$; luminance, 16 cd/m²). The fixation spot and saccade target were black dots 1° in diameter. The test stimulus was a clearly visible vertical green bar, 2° wide and 40° high, presented for a single frame (4 ms) at an intensity nearly three times that of the red background. The contrast of the stimulus was always one log unit above detection throughout the entire course of the saccade (Diamond et al. 2000). For the data reported here, a black ruler was present on the screen 3° below the level of the fixation point. However, all conditions were also performed with the ruler presented 1000 ms after saccadic onset, giving substantially similar results as has been observed previously (Morrone et al. 1997; Ross et al. 1997).

Eye movement measurement: Eye movements were monitored by an infrared limbus eye tracker (HVS SP150) with 1000 Hz temporal resolution. The horizontal resolution was 0.01 degree, and accuracy was 0.1 degree (manufacturer's specifications). In practice we observed an accuracy of about 0.2 degrees over the range we studied. The infrared sensor was mounted below the right eye on transparent wraparound plastic goggles through which subjects viewed the display screen binocularly. The sensor was mounted close to the eye, so the subjects were unaware of its presence. The head of the subject was constrained by a chin-rest so that the eye was 37 cm from the screen. The PC sampled eye position at 1000 Hz and stored the trace in digital form after suitable linearization. Before each session, both the gain and the linearity of the eye tracker were calibrated by asking subjects to make saccades to five fixed points arranged horizontally. Calibration was checked frequently during the recording sessions. Eye traces of subjects, together with target and stimulus presentation times, calibration, and subject responses were stored in digital form for later off-line analysis.

Procedure. Observations were made in a dimly lit room, where the frame of the monitor was clearly visible. There were 3 types of trials:

1. *Unadapted trials.* These were similar to those reported in a previous experiment (Morrone et al. 1997). Trials began with a dark fixation spot, which appeared usually 14° to the left of the center of the screen and stayed on thereafter. After a warning the fixation spot disappeared and at the same time a target appeared (and thereafter stayed on) at center (0°), 14° right of center, or 20° right of center; subjects immediately saccaded to it, producing 14, 28 or 34° saccades. On each trial a green bar was briefly presented (for one frame: 4 ms) either before, during, or after the saccade. Subjects were required to report the positions of bars with reference to the ruler by calling out a number to the experimenter. Subjects tended to report positions in integral numbers of degrees. If s/he did not see it, that trial was annulled (some conditions were particularly difficult to see, particularly bars in the far periphery, sometimes resulting in a paucity of data at a particular region). The arrangement of fixation spot, saccade target, ruler, and bar are shown in the insert of Figure 1 (ignoring the white points at present). The experimenter checked the quality of each saccade and the computer's estimate of saccade onset, and recorded the subject's reported position on the computer.

Average latencies for each subject were estimated after the first few trials, and this information was used to attempt to display the bar at a given time with respect to saccadic onset. However, actual delays were calculated off-line.

2. *Adapted trials:* During saccadic adaptation, the target was caused to step as soon as the computer detected the onset of a saccade. When the horizontal eye position signal reached a pre-determined threshold (relative to fixation), the target shifted abruptly to a new position, either closer to (gain decreasing) or further from (gain increasing) the initial target. After an initial training session (see later), the saccade amplitude of the subject adapted gradually (usually within 20 trials) so the endpoint of the first saccade became progressively nearer to the new position of the target (Figure 1A). We used three different intrasaccadic target steps: from 14° to 20° , producing an increase in saccadic gain; from 14° to 0° , producing a decrease in saccadic gain; and from 8° to 14° , producing an increase in saccadic gain (while not encroaching on the monitor border). As before, subjects were asked to report the position of the bar flashed before, during, or after the saccade, but the response was used only if the

adaptation was within 5% of the actual target displacement.

Types of trials were blocked within a given session (gain-increasing, gain-decreasing etc), with 200-300 trials per session. The starting and finishing point of the trials was kept constant within each session, to produce strong and stable adaptation, given the position specificity of adaptation (Alahyane and Pelisson 2003, 2004). Subjects usually perceived the target to be at the postsaccadic center of gaze. During the adaptation process the subjects were sometimes aware of their own saccadic inaccuracy as evidenced by the necessity of the compensatory saccade. Fully adapted subjects were generally unaware of the adaptation process. The subjects never noticed the intrasaccadic shift. We assume that subjects did not use a deliberate strategy of gain change, as it did not occur immediately or all at once after a delay, but gradually over time (fig 1B). During conditions of adaptation a trial was considered valid if the saccadic amplitude was within 5% of the actual displacement to the final target, and if the target had been stepped to the final position within 10 ms of saccadic onset.

3. Control trials. We used two different types of control trials for some of the adaptation studies. In one we occasionally (on 20% of trials) interposed trials in which the subjects were asked to judge the position of the bar without making any saccade at all (to see if the perceptual effects of saccadic adaptation were a general result of the adaptive process or required an actual saccade). Here the fixation spot remained on and the targets did not appear. The trial was only accepted if the subject continued fixation throughout the trial.

In another control, we interposed 15% *catch trials* in the adapted state experiments, in which the target never stepped on saccade trigger, but just disappeared. In these trials the subjects were still asked to report the position of the bar (to see if the intrasaccadic target step caused any perceptual effect, or if the subjects used the final target position as an aid to localize the flash.).

Subjects. Full data in the major conditions was collected from four subjects, two authors and two naïve to the goals of the experiment. However, all authors served as preliminary subjects for all conditions, and their results agreed with those of the more thoroughly studied subjects. We took particular care to randomize both the position of the bars and the time of their

presentation within a given experimental condition, so subjects (both authors and naïve) were completely unaware of the conditions of a particular trial, given that people are very poor in localizing the time of presentation of a brief stimulus around the time of the saccade (Ross et al. 1997; Schlag and Schlag-Rey 2002). This minimizes the possibility of response stereotyping or other forms of bias in the data. The use of two naïve subjects for most major conditions further excludes this possibility.

Before collecting data on the adaptation conditions, we trained subjects were for one or two days on the various adaptation regimes, until adaptation occurred within the first 20-40 trials. For each adaptation condition 1000 trials were typically collected. Considering all the trials performed, including training trials, a small number of trials did not meet our stringent criteria of adaptation and were therefore eliminated. In many of the discarded trials the subjects made compensatory second saccades. However, most of these errors occurred early in the adaptation process. Restricting the analysis to the session after training to achieve good adaptation, on average 83 percent of all trials could be used for off-line analysis. Adding the remaining saccades did not change the overall pattern of results.

The experimental protocol was approved by the Consiglio Nazionale di Ricerca (CNR) as being within its regulations for human experimentation, under the Helsinki protocol.

RESULTS

Adaptation of saccadic gain

Before measuring visual mislocalizations around the time of saccades, we trained subjects in the saccadic adaptation paradigm of McLaughlin (1967). Subjects viewed a dark fixation spot 14° to the left of the center of the screen (see fig 1C). After a warning a target appeared 14° right of center, to which subjects immediately saccaded. As soon as the saccade began, the target shifted to the new position (Figure 1A). Initially, subjects made two separate saccades, first to the initial saccadic target followed by a corrective saccade to the new position (dark trace of Figure 1A). Over the course of about 20 trials, the amplitude of the first saccade changed so that the endpoint of the saccade approximated the new position of the target, with

the second saccade become smaller and of decreased latency (Figure 1A). During this time subjects ceased to perceive the intrasaccadic target shifts, and usually perceived the target to be at the postsaccadic center of gaze.

The effect of saccadic adaptation on peri-saccadic spatial perception

This experiment was basically like that of Ross et al. (1997), except we also measured localization under conditions of gain-increasing and gain-decreasing adaptation. Bright green bars were briefly flashed to one of four positions (-4° , 4° , 8° and 18°), and subjects were required to report where they seemed to appear (see figure 1 and *methods* section for more details). In all conditions reported here, both the time of presentation relative to the target presentation and the actual spatial position was completely randomized, so subjects were unaware of the spatial and temporal condition of any given trial. The actual time of stimulus presentation relative to saccadic onset was determined during offline analysis. During this analysis we only accepted trials in which the saccade was within 5% of the final target (although data collection commenced before adaptation was complete).

Figure 2 shows average perceptual localization as a function of time relative to saccadic onset, for one of the four subjects used in this study (an author in this case). The three panels on the left show data for normal unadapted saccades of amplitude 14° , 28° and 34° , starting from fixation at -14° . Within each panel the mean and standard error of the reported position of the bar is shown as a function of presentation time relative to the saccade; the horizontal lines of matched color show the actual position of the bars (-4° : red, 4° : blue, 8° : green and 18° : black). In this unadapted condition, the subject reported the bar near its veridical position when it is presented long before or long after the saccade. When the bar was presented in the interval immediately before or during the saccade, however, the subject made gross errors of mislocalization, reporting the bars close to the saccade target (T). The mislocalization errors were maximal near saccadic onset, and fell off steeply in a well-ordered fashion before and after, as described in detail elsewhere (Lappe et al. 2000; Morrone et al. 1997; Ross et al. 1997; Ross et al. 2001).

The two panels on the right show results for conditions of adaptation, either gain decrease of 50% (upper) or gain increase of 21% (lower), both with initial saccadic targets 28° to the right of fixation. Subjects had no difficulty in localizing the bars in the adapted

conditions, and reported that the task was subjectively very similar to that in the unadapted condition. Again there was very strong peri-saccadic mislocalization, leading to a transient compression of space. The focus of the compression was near the actual saccade goal rather than to the original saccade target. As for unadapted saccades, presenting the ruler 1 second after the stimulus had been displayed, rather than continuously, did not affect the results.

Furthermore, and quite unexpectedly, there was a systematic distortion of perceived position in some regions during the entire pre-saccadic fixation interval. In the gain decrease experiment, positions 4° and 8° (blue and green symbols) were severely mislocalized by as much as 8° in the direction of the gain decrease. This occurred very early with respect to saccadic onset, sometimes even before the appearance of the saccadic target (and the disappearance of the fixation point). Note that this distortion does not follow the same pattern as the peri-saccadic compression towards the saccadic target. In this case the shift is in the direction of the adaptation, although not uniform for all spatial positions.

For the gain increase experiment the major early mislocalization occurred for position 18° (black symbols). Here the mislocalization was again strong, up to 6° , but in the other direction, again in the direction of the saccadic adaptation. In both cases the region of early mislocalization includes the region between the initial and stepped target positions. This result implies a form of trans-saccadic distortion limited to an area around the saccade target and its ultimate goal.

To rule out the possibility that the compression to the final target was caused by the physical presence of that target, we performed another control experiment using “catch trials”. Every now and again (unpredictably, but on average 1 trial in 5) we presented well-adapted subjects with trials in which the saccade target disappeared but never reappeared. The subjects made the adapted saccades as expected even though there was no intrasaccadic target step. In these conditions, as before, compression was towards the final landing point of the eye, even though there was no actual target there (Figure 3). The early mislocalization in the region between saccadic target and landing point also occurred.

We studied peri-saccadic localization (for trials that occur in the 20 ms interval prior to saccadic onset) over a finer spatial scale. The results of four subjects (two authors and two naïve) are shown in figure 4, as a function of real spatial position of the bars. The adapted (filled symbols) results are shown together with unadapted saccades of the same actual amplitude (open symbols). The results of all subjects are quite similar in both conditions.

Compression tended to be around the final landing point of the eye, not the intended eye-movement. That is to say, irrespective of the actual position of where the bar was presented, it tended to be seen at the second saccadic target, 0° in the case of the gain-decreased saccades and 20° in the case of the gain-increased saccades. There were some individual differences in this general trend, particularly for subject MCM who showed less compression for the more peripheral targets with the gain-decreased saccades.

To describe better the effects of adaptation on saccadic compression, we calculated the “focus of the compression” for all four subjects. The focus was calculated from the data of figure 4, by regressing the errors in localization against real position, and estimating the zero intercept. These estimates are shown in figure 5 for all four subjects. The abscissa indicates the five relevant conditions: short, medium and long unadapted saccades, together with gain decreasing and gain increasing adapted saccades, where the first target corresponded to the medium saccade, and the final target to the short or long saccade respectively. For the normal saccades, the compression was usually very close to the saccadic goal, at 0 , 14 or 20° . With adaptation, the compression was more towards the final goal than the initial target. There were some variation between subjects, but the average trend (large gray symbols) was clearly towards the saccadic endpoint.

A one way ANOVA revealed a highly significant effect of saccade type ($p < 0.001$, $F=66$, $DF=4$). Pair-wise multiple comparisons of the group data revealed significant differences between all possible pairs **except** between normal saccades to 0 and gain-decreasing saccades with endpoint 0° ($p=0.5$), and between normal saccades to 20° and gain-increasing saccades with endpoint 20° ($p=1$). The average compression foci for both adaptation conditions were significantly different from those to the primary saccade target corresponding to the intention to move (14°), $p < 0.001$ for gain-decreasing and $p=0.02$ for gain-increasing. This confirms the suggestion that the focus of saccadic compression during adaptation tends to be the final landing point of the eye, rather than the initial target that elicited the eye-movement.

A possible problem with the gain-increased saccades was the effect of the screen border. To minimize this factor, we studied an adaptation sequence in which the initial target point was closer to the center of the screen (8° right of center), so the second point remained quite far from the edge (16° from the edge). Here also compression moved to the actual endpoint of the saccades, whether they were adapted or unadapted (Figure 6).

The extent of trans-saccadic perceptual distortion

As noted in figure 2, stimuli presented to some spatial positions well before the saccade were also mislocalized, even when presented several hundred milliseconds before saccadic onset. To examine this phenomenon further and to determine its spatial extent, we presented bars well before a saccade (more than 150 ms) to all spatial positions. Again the spatial positions and presentation times were randomized from trial to trial to minimize response stereotyping.

Figure 7 shows the results for same four subjects of figure 4, two authors (above) and two naïve subjects unaware of the goal of the experiment (below). The data on the left show results for adapted saccades, both gain-increased (open circles) and gain-decreased (filled squares), with the dotted line showing veridical response. The deviations from this line (errors in mislocalization) are not uniform over space, but extend roughly from the fixation point towards the saccade goal. They are strongest in the region that lies between the saccade target and its ultimate landing point. There was also distortion in the region immediately adjacent to this, and also some variation from subject to subject, but all subjects showed a robust shift in the direction of the adaptation, and only over a limited area centered around the region between saccadic targets.

For unadapted saccades, localization of objects flashed well before saccadic onset showed only a small deviation from the veridical, usually in the same direction for the short and long saccades (Figure 7, right). A two way ANOVA was performed for each subject, revealing significant differences ($p < 0.05$ in all cases) between gain-increasing and gain-decreasing adapted conditions, but no significant differences for localization during normal saccades.

Given that the distortions during saccadic adaptation occur for very large durations prior to saccade onset, we asked if the saccade were necessary at all: perhaps space becomes distorted during adaptation, even for normal viewing. To test this we interposed trials in which the fixation point never disappeared, and instructed subjects not to make a saccade in these conditions. For these trials, shown in figure 8, the pattern of results was quite different from for the trans-saccadic localization. There were some small deviations from veridical (especially for subject HA) but these were much less than with the saccade, and the shift was not specific for the direction of the adaptation.

DISCUSSION

These experiments report two main effects of postsaccadic spatial localization during saccadic adaptation. Firstly, targets briefly presented around the time of saccades show the type of compressive mislocalization pattern previously reported for normal saccades (Ross et al. 1997). The focus of the compression tends to be closer to the endpoint of the actual movement, rather than the target that evoked the saccade, and indeed the entire mislocation curves of adapted saccades are very similar to those obtained with unadapted saccades of the same amplitude. Secondly, during periods of adaptation, stimuli presented well before the saccade to certain regions of space were mislocalized

That the compression focus should follow the actual rather than the planned saccade is not entirely expected from previous work on adaptation. For example, during saccadic adaptation, the attentional focus of human subjects, measured by contrast sensitivity, remains at the spatial location of the original target, even though the actual saccade is made to a different target (Ditterich et al. 2000). In those experiments subjects had to report the orientation of an object but not its position. Our experiments did not examine stimulus detectability or other indices of attention, but localization of a high contrast and clearly visible bar. It is interesting that although the attentional focus remains with the initial saccadic target, saccadic mislocalization occurs towards the endpoint of the actual movement.

Compression did not depend on postsaccadic visual information from the target itself. We performed controls where the subject was well-adapted but rather than stepping to a new position the target disappeared completely. The results with these catch trials were essentially the same as when the target was moved, showing that the actual presence of the target was not necessary as a visual or attentive reference in these experiments. However, in the absence of a postsaccadic target, memory information about the usual postsaccadic target position in the adaptation trials might have been used together with physical visual references provided by the ruler and the monitor edges (Lappe et al. 2000). The role of visual references in peri-saccadic mislocalization has been debated over several years. Although it is clear that visual references have a strong effect on mislocalization (e.g. Honda 1999) and, in particular, on compression (Lappe et al. 2000), mislocalization does occur in total darkness (eg Cai et al. 1997; Matin and Pearce 1965). Recently it has been shown that visual compression also

occurs in darkness, provided that the mislocalization of the saccadic target itself is considered (Awater et al. 2000). These results show that compression does not result entirely from visual references, but may also involve an extra-retinal signal.

There are two general possibilities how an an extraretinal signal could act in transsaccadic spatial localization. An extraretinal signal could be used to construct an impression of position of a visual target in space (Matin, 1965) but this role is only clearly applicable when targets are in darkness. When other visual references are available the role of the extraretinal signal becomes more complex (cf Honda 1999, Dassonville et al., 1995, Deubel et al., 1998). For example, an extraretinal signal could affect mechanisms that determine relative visual position (Lappe et al., 2001). The *reference object theory* of transsaccadic spatial localization (Deubel et al., 1998) assumes that the position of objects before the saccade is encoded relative to reference objects, the most obvious reference being the saccade target. After the saccade, the position of the object has to reconstructed from the encoded relative position, the extraretinal signal, and the reference position of the saccade target. Lappe et al. (2001) proposed that the compression originates in the encoded relative distance between flash and target.

How would the data of our experiments fit with the reference object theory? One would have to assume that the flash position is first encoded - and compressed - towards the initial visual cue at T1, and after the saccade reconstructed with respect to the postsaccadic target position at T2, using a combination of visual and extraretinal references on T2's position itself. Thus, the center of compression should eventually lie at T2. However, the theory essentially predicts that this is the result of two separate processes: perisaccadic compression of relative distances in the scene encoded at the time of the flash plus a transformation from T1 to T2 because of the saccade adaptation. The long-lasting space distortion that we observed long before the saccade can help to distinguish these two stages because it clearly relates only to the transformation induced by saccade adaptation. The theory predicts that perisaccadic effects can be seen in isolation when the adaptation effects are subtracted. For instance for the data in Figs. 2 and 3 this would mean to shift each curve such that the data points long before the saccade are aligned with the true positions (dotted lines). Indeed, the perisaccadic data points of the blue, green, and black curves in the Figs. 2 and 3 would then fall close to position T1, consistent with the theory. The red curves, however, do not follow the prediction.

The functional role of saccadic compression is still poorly understood, although it has been confirmed in many laboratories over several years. However, whatever its actual role, it is a useful tool to study peri-saccadic mislocalizations, as they can be characterized by a single attractor point. The fact that peri-saccadic mislocalizations during adaptation follow closely the pattern generated by an unadapted saccade of same actual size can be interpreted in one of two ways: either there exists an efference copy signal of the same size as the actual saccade, or the effect results entirely from the post-positioning of the eye. The second possibility would suggest that visual stimuli briefly flashed around the time of the saccade are always seen at the final landing point, irrespective of whether the saccade was planned to that point, or whether the landing point was different from that expected, due to adaptation. Perhaps objects presented around the time of saccades are not localized well by the visual system, because of demonstrated suppression of the magno-cellular system (Burr et al. 1994; Ross et al. 1996). Indeed, the strength of compression correlates with the strength of suppression (Michels and Lappe, 2004). The default localization of all stimuli flashed around the time of the saccade is the fovea. In any event, this explanation also requires an extra-retinal corollary discharge suppression signal at the time of the saccade (Diamond et al. 2000). A somewhat similar explanation was proposed by Niemeier et al (2003), who showed that saccadic suppression of displacement occurred most dramatically at the saccade goal, and correlated with the unreliability of individual subjects' saccades. They interpreted these data to suggest that the brain uses an optimal inference solution one property of which is to compress stimuli towards the center of gaze.

The alternate explanation is that the mislocation is driven by an efference copy signal matched to the actual not intended saccadic amplitude. It is not obvious where this extra-retinal signal may come from, particularly in the adapted condition. Saccadic adaptation, although driven by visual error, occurs on the motor side, involving the fastigial nucleus (Robinson and Fuchs 2001) and vermis (Barash et al. 1999). PET studies in humans implicate the cerebellum (Desmurget et al. 1998). Neurons in oculomotor supranuclear areas such as the superior colliculus do not show evidence of adaptation (Frens and Van Opstal 1997), suggesting that the colliculus and the cortical areas that project to it specify target location in unadapted visual coordinates, and that the motor program for the adapted saccade is determined by downstream centers. However, the peri-saccadic localization errors seemed to be driven by the final landing point in the adapted condition, suggesting that if an efference copy signal is driving this compression, the signal must arise after cerebellar

modification of the visually determined saccade goal. This signal would then have to ascend to the visual areas of the cerebral cortex. Recently, Sommer and Wurtz (2002) showed that a lesion in the medial dorsal nucleus of the thalamus, which relays the superior colliculus saccadic signal to the frontal eye field, partially affects the monkey's ability to perform double-step saccades, a task for which efference copy is necessary. This pathway, which would be expected to be unaffected by saccadic adaptation, could not supply the entire efference copy signal in the adapted state.

Besides the data on peri-saccadic localization, showing a dynamic remapping at the time of the saccade, we also showed a long-lasting distortion of spatial localization of targets presented long before the saccade. This result bears on issues of spatial memory and trans-saccadic integration. This result bears some resemblance to that of Bahcall and Kowler (1999), who showed spatial mislocalization of objects presented in darkness near the locus of the saccade target. They limited their investigation to perceptual localization of the first saccadic goal, and concluded that the efference copy that compensated for the physical motion of the image across the retina was driven by the "intention to move" signal rather than the actual eye movement. The two studies agree in demonstrating perceptual mislocalizations of stimuli displayed well before saccade onset, in the direction of the adaptation. We further showed that the mislocalization occurred over a limited area of the visual field, the area near the saccade goal and the final saccade endpoint. For the rest of the visual field postsaccadic estimation of the spatial location of a pre-saccadic stimulus is accurate except for times very near saccadic onset.

Bahcall and Kowler explained their result in terms of an inappropriate efference copy, set by the intention to move signal rather than the actual motion. While this explanation could explain our results near the saccadic target, it cannot explain the whole pattern of results shown in figure 7: a single inappropriate efference copy signal would shift the entire curves leftward or rightward by a constant amount, proportional to the mismatch. But our results clearly show that the distortion of space is limited to a region including the first and second saccadic targets. Bars displayed outside this region were localized veridically. For this veridical region the mechanism that compensates for saccades must be an accurate efference copy of the actual, not the intended saccade, or some other signal. There must be a special mechanism that compensates for the part of space around the saccadic targets. This part of space, which is distorted by the motor system, seems to be also distorted by the perceptual

system for a long interval between the appearance of the target and the performance of the saccade.

The pre-saccadic long-lived shift requires that the subject actually make a saccade and report the location of the flashed object after the saccade. Our subjects did not show such a shift when, even though they were adapted, they reported the location of a flashed object without performing a saccade. Moidell and Bedell (1988) reported a small shift (0.5°) in a similar saccadic adaptation experiment without actual saccades. However, this shift is of much lower magnitude than that reported here, below the resolution of our experimental technique.

One exciting possibility is that the pre-saccadic distortion that we observe here constitutes, or at least contributes to, the actual mechanism of saccadic adaptation. However, this tempting idea seems unlikely for many reasons. Firstly, a great deal of evidence suggests that saccadic adaptation occurs in the motor system rather than the visual system (eg Wallman and Fuchs 1998). Secondly, although saccadic adaptation does show spatial selectivity (Deubel 1991; Frens and van Opstal 1994; Noto et al. 1999), the reported spatial selectivity of the adaptation field is not co-extensive with the area of long-lived perceptual shift. In gain-decreasing adaptation, the area beyond the target position is strongly adapted, yet we found little or no long-lived perceptual shift in this region. Thirdly, we found this mislocalization only for briefly presented targets. We have no information of whether targets continuously displayed for longer periods would be mislocalized. Therefore, although we cannot exclude completely this hypothesis, it seems likely that the mechanisms underlying the parametric adaptation field and the long-lived perceptual shift are not the same.

The results for pre-saccadic localization may also be considered in terms of the previously mentioned “*reference object theory*” (Deubel 2003; Deubel et al. 1998), that assumes that pre- and post-saccadic “snapshots” are linked by fixing perceptually the saccadic target, and a limited region around it, that the visual system assumes to be stable. In our experiment, the saccadic target stepped imperceptibly during the saccade so the post-saccadic fixation should be linked to a different portion of the pre-saccadic fixation, causing a local distortion. This distortion was limited to a region of about 20° around the saccadic targets. With a quite different experimental paradigm, Deubel (2003) showed distortion limited to about 3° of space: but in his experiment the saccades were only 6° , compared with the $28\text{-}34^\circ$ saccades used here, suggesting that the extent of the region depends upon saccadic

amplitude.

These data reinforce the concept that human spatial memory, introspectively robust, is in fact rather fragile. It depends on a number of independent metrics that are ordinarily in register, but which can be dissociated. When eye movements and the visual scene are out of joint, the brain struggles as best it can, using both corollary discharge and memory. On the other hand, these data also demonstrate a high potential for plasticity in perception, limited to the relevant region of visual space.

Acknowledgments

This research was supported by grants from the Human Frontiers Science Program, the W. M. Keck Foundation, the National Eye Institute (1 R01 EY014978-01), and the Italian Ministry of Education and Research (MUIR Cofin).

REFERENCES

1. **Alahyane N and Pelisson D.** Adaptation of saccadic eye movements: transfer and specificity. *Ann N Y Acad Sci* 1004: 69-77, 2003.
2. **Alahyane N and Pelisson D.** Eye position specificity of saccadic adaptation. *Invest Ophthalmol Vis Sci* 45: 123-130, 2004.
3. **Awater H, Burr DC, Goldberg ME, Lappe M and Morrone MC.** Peri-saccadic localization after saccadic adaptation. *Perception* 30 (suppl): 24, 2001.
4. **Awater H, Krekelberg B and Lappe M.** Perisaccadic visual stimulation changes perceived target positions. *Perception* 29: 8, 2000.
5. **Bahcall DO and Kowler E.** Illusory shifts in visual direction accompany adaptation of saccadic eye movements. *Nature* 400: 864-866, 1999.

6. **Barash S, Melikyan A, Sivakov A, Zhang M, Glickstein M and Thier P.** Saccadic dysmetria and adaptation after lesions of the cerebellar cortex. *J Neurosci* 19: 10931-10939, 1999.
7. **Bridgeman B, Hendry D and Stark L.** Failure to detect displacement of the visual world during saccadic eye movements. *Vision Res* 15: 719-722, 1975.
8. **Burr DC, Morrone MC and Ross J.** Selective suppression of the magnocellular visual pathway during saccadic eye movements. *Nature* 371: 511-513, 1994.
9. **Cai RH, Pouget A, Schlag-Rey M and Schlag J.** Perceived geometrical relationships affected by eye-movement signals. *Nature* 386: 601-604, 1997.
10. **Dassonville P, Schlag J and Schlag-Rey M.** Oculomotor localization relies on a damped representation of saccadic eye displacement in human and nonhuman primates. *Vis Neurosci* 9: 261-269, 1992.
11. **Dassonville P, Schlag J and Schlag-Rey M.** The use of egocentric and exocentric location cues in saccadic programming. *Vision Res* 35: 2191-2199, 1995.
12. **Desmurget M, Pelisson D, Urquizar C, Prablanc C, Alexander GE and Grafton ST.** Functional anatomy of saccadic adaptation in humans. *Nat Neurosci* 1: 524-528, 1998.
13. **Deubel H.** Adaptive control of saccade metrics. In: *Presbyopia Research*, edited by Stark GOaLW. New York: Plenum Press, 1991, p. 93-100.
14. **Deubel H.** Localization of targets across saccades: role of landmark objects. *Visual Cognition*: 11:173-202, 2003.
15. **Deubel H, Bridgeman B and Schneider WX.** Immediate post-saccadic information mediates space constancy. *Vision Res* 38: 3147-3159, 1998.
16. **Deubel H and Schneider WX.** Saccade target selection and object recognition: evidence for a common attentional mechanism. *Vision Res* 36: 1827-1837, 1996.

17. **Diamond MR, Ross J and Morrone MC.** Extraretinal control of saccadic suppression. *J Neurosci* 20: 3449-3455, 2000.
18. **Ditterich J, Eggert T and Straube A.** Relation between the metrics of the presaccadic attention shift and of the saccade before and after saccadic adaptation. *J Neurophysiol* 84: 1809-1813, 2000.
19. **Frens MA and van Opstal AJ.** Transfer of short-term adaptation in human saccadic eye movements. *Exp Brain Res* 100: 293-306, 1994.
20. **Frens MA and Van Opstal AJ.** Monkey superior colliculus activity during short-term saccadic adaptation. *Brain Res Bull* 43: 473-483, 1997.
21. **Honda H.** Perceptual localization of visual stimuli flashed during saccades. *Percept Psychophys* 45: 162-174, 1989.
22. **Honda H.** Modification of saccade-contingent visual mislocalization by the presence of a visual frame of reference. *Vision Res* 39: 51-57, 1999.
23. **Hopp JJ and Fuchs AF.** Investigating the site of human saccadic adaptation with express and targeting saccades. *Exp Brain Res* 144: 538-548, 2002.
24. **Kommerell G, Olivier D and Theopold H.** Adaptive programming of phasic and tonic components in saccadic eye movements. Investigation in patients with abducens palsy. *Invest. Ophthalmol.* 15: 657-660, 1976.
25. **Lappe M, Awater H and Krekelberg B.** Postsaccadic visual references generate presaccadic compression of space. *Nature* 403: 892-895, 2000.
26. **Matin L and Pearce DG.** Visual perception of direction for stimuli flashed during voluntary saccadic eye movements. *Science* 148: 1485-1488, 1965.
27. **McLaughlin SC.** Parametric adjustment in saccadic eye movements. *Percept. Psychophys.* 2: 359-362, 1967.

28. **Michels L and Lappe M.** Dependence of saccadic compression and suppression on contrast. *Vision Res.* 44: 2327-2336, 2004.
29. **Moidell BG and Bedell HE.** Changes in oculocentric visual direction induced by the recalibration of saccades. *Vision Res* 28: 329-336, 1988.
30. **Morrone MC, Ross J and Burr DC.** Apparent position of visual targets during real and simulated saccadic eye movements. *J Neurosci* 17: 7941-7953, 1997.
31. **Niemeier M, Crawford JD and Tweed DB.** Optimal transsaccadic integration explains distorted spatial perception. *Nature* 422: 76-80, 2003.
32. **Noto CT, Watanabe S and Fuchs AF.** Characteristics of simian adaptation fields produced by behavioral changes in saccade size and direction. *J Neurophysiol* 81: 2798-2813, 1999.
33. **Robinson FR and Fuchs AF.** The role of the cerebellum in voluntary eye movements. *Annu Rev Neurosci* 24: 981-1004, 2001.
34. **Ross J, Burr DC and Morrone MC.** Suppression of the Magnocellular Pathways during Saccades. *Behavioural Brain Research* 80: 1-8, 1996.
35. **Ross J, Morrone MC and Burr DC.** Compression of visual space before saccades. *Nature* 386: 598-601, 1997.
36. **Ross J, Morrone MC, Goldberg ME and Burr DC.** Changes in visual perception at the time of saccades. *Trends Neurosci* 24: 113-121, 2001.
37. **Schlag J and Schlag-Rey M.** Through the eye, slowly: delays and localization errors in the visual system. *Nat Rev Neurosci* 3: 191-215, 2002.
38. **Sommer MA and Wurtz RH.** A pathway in primate brain for internal monitoring of movements. *Science* 296: 1480-1482, 2002.
39. **Wallman J and Fuchs AF.** Saccadic gain modification: visual error drives motor adaptation. *J Neurophysiol* 80: 2405-2416, 1998.

Revision of 17 December 2004




FIGURE 1

A. The time course of the various events that occurred on each trial in a particular adaptation condition (gain decreasing). The trial starts with the observer looking at the fixation point at left (position -14°). At an unpredictable moment, the fixation point disappears and the target (T1: position $+14^\circ$ in this example) appears at a particular position to the right, to which the observer made a saccade as rapidly as possible. During adaptation trials, the target was moved to T2 as soon as the computer had detected the onset of a saccade. The black trace shows a saccade early in the adaptation sequence, where the observer made two distinct saccades. The gray trace shows a later, perfectly adapted, saccade that goes straight to the final destination.

B. Gain of the saccades (ratio of actual movement to distance to first target) in one experimental session, after the training sessions. After a few trials the saccade goes straight to the final goal, half the initial distance (gain = 0.5). The curve passing through the data is an exponentially decaying fit with a constant of 19 trials. Even after adaptation, not all the trails met the strict criterion of being within 5% of the amplitude to T2.

C. Illustration of the stimulus conditions for the experiments. Note that the fixation (-14°), saccadic targets (0° and 14°) and bar were never seen simultaneously, but followed the time course reported in figure A.




FIGURE 2

Average perceived position of bars flashed at various times relative to saccadic onset (0 ms), in various positions under various conditions, for 1 subject HA. Each point represents the average of 4-10 measurements, with error bars indicating standard error of the mean. Each point is positioned at the mean time of bar appearance for the measurements. For each type of saccade, the bars were flashed at one of four possible positions (randomly interspersed) indicated by different colors: -4° (red), 4° (blue), 8° (green) and 18° (black). The horizontal lines indicate the actual position and the symbols the average perceived position, together with the standard error. The three curves on the left show normal unadapted saccades, of amplitude 14, 28 or 34° illustrated by the vertical arrows at left. The curves on the right show results for gain decreasing and gain increasing saccades, illustrated by the arrows and targets at right (first target T1 in gray). In unadapted saccades, the subjects grossly mislocalized all

bars towards the saccadic target during the period near saccadic onset. In both the gain increasing and gain decreasing condition, they tended to be seen at the final landing point of the eye, not the initial target that elicited the saccades. In both adapted conditions, there was also a strong mislocalization during the entire period prior to saccadic onset for bars flashed in the region between the two saccadic targets. The mislocalization was in the direction of the adaptation.




FIGURE 3

“Catch trials”: after an initial period of adaptation, on every third to seventh trial (1/5 on average) the first target T1 disappeared after the saccade but the second target T2 did **not** appear in its new position on that particular trial. Each data point is the average of 2-4 measurements, with bars representing standard error. On the other trials (not reported here) the normal adaptation sequence (with target displacement) continued to maintain adaptation. Although there are necessarily far fewer trials in this condition, the results basically confirm those of figure 2, showing that it is not the presence of the second target that causes the mislocalization.




FIGURE 4

Mislocalization during the interval just prior to saccadic onset (-20 – 5 ms), as a function of space, for gain decreasing (left) and gain increasing (right) saccades. For comparison unadapted saccades of 28° amplitude (matched to actual movement) are shown as open triangles. Each point represents the average of 4-10 measurements, with error bars indicating standard error of the mean. The dotted lines indicate the expected results for veridical perception, and the symbols the average perceived position (average of 4 – 10 trials), together with error bars. The arrows indicate the fixation point (lowest arrow in each panel), the first saccadic target (pale grey) and the second saccadic target. In all cases there were gross distortions in perceived position, with a strong compression around the saccadic target. In general, the compression was to the final landing point of the eye (dark arrows), not to the initial target (light arrows) that elicited the saccades. For the gain decreasing saccades for MCM, the compression was limited to the 14° of space between fixation and landing point.




FIGURE 5

Focus of compression for the four subjects for whom complete data was taken. The focus was calculated from the data of figure 4, regressing the errors (deviations from real position) against real position and estimating the intercept, weighting each point by its standard error. If the linear regression for all spatial positions accounted for less than 80% of the variance, the range of the regression was limited to points to the right of fixation. The time window was between -20 and +5 ms relative to saccade onset. The filled data points show foci for individual observers, with error bars indicating standard error of the intercept estimation. The open data points are the average results, with error bars indicating standard error between subjects. A one-way ANOVA revealed a highly significant main effect ($p < 0.001$, $F = 66$, $DF = 4$). Pair-wise multiple comparisons revealed significant differences between all possible pairs except those indicated by the arrow labeled “NS”. The average compression foci for both adaptation conditions were significantly different from those to the primary saccade target (14°), $p < 0.001$ for gain-decreasing and $p = 0.02$ for gain-increasing. HA and CM were authors; SD and RG were naïve.




FIGURE 6

Gain increasing saccades of smaller amplitude ($22 \rightarrow 28^\circ$), finishing well away (16°) from the edge of the screen, to exclude the problem of visual interference of the edge. Fixation was at -14° , T1 at $+8^\circ$ and T2 at $+14^\circ$. The results are substantially the same as those observed with the larger saccades, both for presaccadic presentations (Fig. 7) and peri-saccadic presentations (Fig. 4). Other details are given in the captions to those figures.





FIGURE 7

Perceived position of bars flashed well before saccadic onset (-500 – -150 ms), plotted as a function of real position of the bar. The bars were presented at random to all positions, with a spatial grain of 2°. Each point represents the average of 4-10 measurements, with error bars indicating standard error of the mean. The curves at left show results for adapted saccades (filled squares gain decreasing, open circles gain increasing), and those at right for normal saccades of the same actual amplitude (filled triangles 14°, open triangles 34°). The arrows on the axes indicate the key saccadic targets. The black arrow at -14° the fixation point, and the gray arrow at +14° the initial saccadic target. The black arrows either side of this are the final targets for gain decreasing and increasing conditions. Over a limited region, corresponding roughly to the area between the two saccadic targets, space was grossly distorted during adaptation, in the direction of the adaptation. There were also some mislocalizations during normal saccades, but they were not so systematic, nor of the same magnitude.

**FIGURE 8**

Measurements of apparent position during fixation while subjects were adapted (decreasing and increasing gain). On 20% of trials (on average), the fixation point did not extinguish, and on these trials the subject did not make a saccade: all else was the same as before. Each point represents the average of 3-6 measurements, with error bars indicating standard error of the mean. Here there were some mislocalizations (especially for HA), but lower magnitude than when an actual saccade was made. Importantly, the errors were in the same direction for the gain decreasing as the gain increasing saccades.